

Case Report

A Rare Case of Herpes Simplex Virus Type 2 Induced Acute Retinal Necrosis

Stephen Sudhakar*, Charanya C**, Ananya A**

*Prof & HOD, **Post Graduate Student, Dept. of Ophthalmology, Chettinad Hospital & Research Institute, Chennai, India.



Dr. K. Stephen Sudhakar did his MBBS from PSG Institute of Medical Sciences, Coimbatore and M.S in Ophthalmology from Madras Medical College, RIOGOH. He completed his DNB from National board. He is a Fellow of International Council of Ophthalmology [U.K]. He completed his Ratan Tata Microsurgical Fellowship from Sankaranetralaya. He has presented papers in many national and international conferences and published articles. He is a life member of All India Ophthalmology Society [AIOS]. Currently working as the HOD and Professor in Department of Ophthalmology. His areas of interest include Neuro- ophthalmology, Ocular traumatology, Medical Retina and Glaucoma.

Corresponding author - Stephen Sudhakar(dr_stephen1978@yahoo.com)

Chettinad Health City Medical Journal 2016; 5(1): 38 - 40

Abstract

29 year old healthy female came with complaints of defective vision in left eye for one week duration along with pain, redness and floaters. On examination, patient was found to have pan uveitis, retinal vasculitis, vitritis and necrosis of the peripheral retina in the left eye. A diagnosis of acute retinal necrosis was made. PCR analysis of aqueous and vitreous revealed Herpes Simplex Virus type 2 which is a rare cause for acute retinal necrosis.

Key Words: Acute Retinal Necrosis, Immunocompetent, Herpes Simplex Type 2

Introduction

Acute retinal necrosis (ARN) is a rare but potentially devastating syndrome characterized by progressive peripheral necrotizing retinitis. It was first described in 1971¹. It is usually caused by Varicella zoster and occurs in immunocompetent individual².

Progressive outer retinal necrosis (PORN) is also type of necrotizing retinitis and characterized by multifocal lesions without granular borders in deep retinal layers³. The main difference between both ARN and PORN is ARN occurs in immunocompetent and PORN in immunocompromised and absence of inflammation in PORN⁴.

Herpes simplex virus is a DNA virus belonging to herpes family is two types – type 1 and type 2. Herpes Simplex Virus 1 is transmitted by oral contact and direct contact. In eye it causes herpes keratitis which is the leading cause of corneal blindness in U.S.A⁵. Herpes Simplex Virus 2 causes genital herpes and sexually transmitted. 3.7 billion people have Herpes Simplex Virus 1 and 417 million people have Herpes Simplex Virus 2 infection⁶.

Case History

A 29 year old female came with complaints of defective vision in left eye for one week duration. There is a history of fever, vomiting and headache for 2 weeks for which she was admitted and treated. There is also history of joint pain. On examination, right eye appears to be normal with normal unaided visual acuity. Left eye anterior segment showed circumciliary congestion of conjunctiva with fine keratic precipitates in cornea and anterior chamber showing cells 3+. Patient's best corrected visual acuity in left eye was 6/12. The patient was given topical steroids and was reviewed after 1 week. Patient vision worsened to 3/60 in spite of

treatment with pupil sluggishly reacting to light. Fundus examination of right eye was within normal limits. Fundus examination of left eye showed hazy media due to vitreous inflammation. Vitreous cells were 2+. Optic disc showed edema with blurring of disc margins(Fig1). The superior retinal quadrant showed pale and thin retina with granular borders which are well demarcated(Fig 2). There was evidence of retinal vasculitis with sheathing of arteries and occluded peripheral retinal vessels in superior quadrant of retina(Fig 3). With the above clinical findings, a presumptive clinical diagnosis of acute retinal necrosis in the left eye was made. Routine blood investigations, ANA, RA factor VDRL, HIV serology, Mantoux, Chest X-ray were within normal limits.




Fig 1: Optic disc edema with vitritis




Fig 2: Thinning and sheathing of retinal arterioles




Fig 3: Necrosed peripheral retina with granular borders

Discussion

Acute Retinal Necrosis is caused by viral infection mostly by Varicella zoster virus or Type 1 Herpes Simplex. Type 2 Herpes Simplex Virus causing Acute Retinal Necrosis is very rare⁷. American Uveitis Society defined ARN based on following clinical criteria⁸:

- 1) One or more foci of retinal necrosis with discrete borders located in the peripheral retina
- 2) Rapid progression in the absence of antiviral therapy
- 3) Circumferential spread
- 4) Evidence of occlusive vasculopathy with arterial involvement
- 5) A prominent inflammatory reaction in the vitreous and anterior chambers

Our patient had all the above clinical findings and hence was diagnosed clinically as Acute Retinal Necrosis. The confirmation of organism is usually by Polymerase Chain Reaction with aqueous or vitreous samples. We went on to do a PCR analysis of aqueous and vitreous and it showed type 2 herpes simplex virus which is very

rare. Rarely in non-responding cases is diagnostic vitrectomy or retinal biopsy needed⁹.

Antiviral therapy can be given systemically as well as intravitreally. Commonly used systemic antiviral used are acyclovir, famciclovir, valacyclovir¹⁰ antiviral agents given intravitreally are ganciclovir (200 microgram/0.1 ml) foscarnet (1.2 mg/0.1ml). They are given in retinitis threatening macula or optic disc¹¹.

The patient completed the course of oral Valacyclovir 1gm TID for 6 weeks and her renal function was stable throughout the treatment. Intravenous anti-viral medications were not given as the patient is immunocompetent and had responded to oral valacyclovir. The patient is currently on regular follow up for recurrence of disease in both eyes.

Conclusion

Acute retinal necrosis is a severe ocular inflammatory syndrome in immunocompetent associated with poor visual outcome. It is commonly caused by Varicella zoster virus, Herpes Simplex virus type 1 and less commonly by Herpes simplex virus type 2 as in our case which was confirmed by PCR analysis of vitreous and aqueous samples. The patient responded well to oral Valacyclovir which was given for 6 weeks along with steroids. Retinal inflammation subsided, but retinal thinning still persisted as the retina was necrosed.

Patient is currently on follow up for recurrent involvement in same eye and for involvement of other eye. Photocoagulation of retina is planned in left eye for prevention of Retinal Detachment.

References

- 1) Urayama A YN, Sasaki T, Nishiyama Y, Watanabe S, Wakusawa S, Satoh Y, Takahashi K, Takei Y. Unilateral acute uveitis with periarteritis and detachment. *Japan J Ophthalmology*. 1971; 25:607-619.
- 2) Young NJ, Bird AC. Bilateral acute retinal necrosis. *Br J Ophthalmol*. 1978; 62:581-590.
- 3) Okunuki Y, Usui Y, Kezuka T, et al. Four cases of bilateral acute retinal necrosis with a long interval after the initial onset. *Br J Ophthalmology*. 2011; 95:1251-1254.
- 4) Cochrane TF SG, McDowell C, Foot B, McAvoy CE. Acute Retinal Necrosis in the United Kingdom: results of a prospective surveillance study. *Eye*. 2012; Eye advance online publication 27 January 2012; doi: 10.1038/eye.2011.338.
- 5) Muthiah MN, Michaelides M, Child CS, Mitchell SM. Acute retinal necrosis: a national population-based study to assess the incidence, methods of diagnosis, treatment strategies and outcomes in the UK. *Br J Ophthalmology*. 2007; 91:1452-1455
- 6) Fisher JP, Lewis ML, Blumenkranz M, et al. The acute retinal necrosis syndrome. Part 1: Clinical manifestations. *Ophthalmology*. 1982; 89:1309-1316.

- 7) Martinez J, Lambert HM, Capone A, et al. Delayed bilateral involvement in the acute retinal necrosis syndrome. *Am J Ophthalmology*. 1992; 113:103-104.
- 8) Holland GN. Standard diagnostic criteria for the acute retinal necrosis syndrome. Executive Committee of the American Uveitis Society. *Am J Ophthalmol*. 1994; 117:663-667.
- 9) Clarkson JG, Blumenkranz MS, Culbertson WW, et al. Retinal detachment following the acute retinal necrosis syndrome. *Ophthalmology*. 1984; 91:1665-1668.
- 10) Lau CH, Missotten T, Salzmann J, Lightman SL. Acute retinal necrosis features, management, and outcomes. *Ophthalmology*. 2007; 114:756-762.
- 11) Fernando A, Esposito GD, Thi Tran, Nathalie Cassoux, Christine Fardeau, PhuLehoang, BahramBodaghi. Long-term Visual Outcome of Patients Presenting With Necrotizing Viral Retinopathies. Association for Research in Vision and Ophthalmology. Fort Lauderdale, FL 2011.

Spirituality Boost in a Nasal Spray

Are you troubled by your lack spirituality? Are you tempted to seek the guidance of a spiritual healer to fill that void? You are more likely to succeed in your goal if you bought a can of Oxytocin nasal spray instead. Oxytocin is a physiological hormone mainly involved in childbirth and breastfeeding. But it is also called "Cuddle hormone or love hormone" as it enhances maternal/social bonding, empathy and sexual pleasure. In a new random controlled study carried out on 83 males, researchers found that Oxytocin, when administered as nasal spray, enhanced the sense of spirituality in the study subjects when compared to those receiving only placebo. The effect was more marked in those subjects who had a particular genetic variant of CD38 gene, which is known to be involved in oxytocin release from hypothalamus. Whether the hormone has similar effect in women is not yet known and needs to be investigated. For the present, those men seeking spiritual high, can achieve that with Oxytocin nasal spray!

(Social Cognitive and Affective Neuroscience, doi: 10.1093/scan/nsw078, published online 17 June 2016)

- Dr. K. Ramesh Rao