

Review Article

Evolution of Stapes Surgery

*P Karthikeyan, **Davis Thomas Pulimoottil

*Professor & HOD, **Senior Resident, Dept. of ENT, Mahatma Gandhi Medical College & Research Institute, Pondicherry.



Currently working as Professor and Head of the Department of Otorhinolaryngology, Mahatma Gandhi Medical College and Research Institute, Pondicherry, Prof. Dr. P. Karthikeyan is a leading ENT surgeon in Pondicherry, with a strong interest in Otolaryngology and Endoscopic Sinus Surgery and its advanced application. He is a keen teacher, with more than 15 years of experience and has received the "Most Popular Teacher" award in 2013, and has been a guide to several postgraduates. Dr. Karthikeyan is also enthusiastic about research in Otorhinolaryngology, and has 14 publications in indexed peer-reviewed journals. He is an active member of the Association of Otorhinolaryngologists of India and is also a reviewer for various national and international journals. He has also successfully organized the 1st Southzone ENT Conference in 2005 and two live surgical workshops on 'Snoring and Sleep Apnoea Surgery' and 'Advanced Otological Surgery'.

Corresponding author - P Karthikeyan (drkarthik73@yahoo.co.in)

Chettinad Health City Medical Journal 2015; 4(4): 167 - 170

Abstract

Otosclerosis is a hereditary localized disease of the otic capsule which is characterized by alternating phases of bone resorption and formation, as a result of which there is removal of mature lamellar bone by osteoclasts and the subsequent replacement by woven bone of greater thickness, cellularity and vascularity in the active stage, and later this is followed by sclerosis and hardening, leading to ankylosis of the stapes foot plate and conductive deafness. The surgical management of this condition has been the subject of much debate and several controversies over the course of more than a century of study, with several stalwarts of the field of Otorhinolaryngology taking opposing points of view in regard to this. Beginning with Kessel in the late 1800s, stapes surgery began with the era of stapes mobilization, which progressed to the era of stapes fenestration following the introduction of the operating microscope into otologic practice, and finally reached the era of stapedectomy in the 1950s. Advances in surgery have come as a result of the introduction of laser, microdrills and now, robotics into the field of stapes surgery and brought us to where we are now. Even still, the basic principles of stapes surgery deduced by the pioneers of the profession still hold true today, albeit with few changes.

Key Words: Stapes surgery, Otosclerosis, Stapedotomy

Introduction

Otosclerosis is a genetically-mediated primary metabolic bone disease of the otic capsule and ossicles, causing ossicular fixation and hence resulting in conductive or mixed hearing loss and is mediated via autosomal dominant transmission with incomplete penetrance and complex, not clearly defined triggering mechanisms¹. It is a disease of young patients, mainly females, who are handicapped by progressive hearing loss for no apparent reason. The diagnosis is usually straight forward and is made on clinical grounds. These patients can be nicely and effectively rehabilitated by modernized hearing aids or surgery, which is the best option in the expert's hands².

Discovery of Otosclerosis

Ludwig van Beethoven (1770-1827) was considered to have mixed hearing loss attributed to otosclerosis. Beethoven first admitted his deafness at the age of thirty-one when writing to a friend complaining of progressive worsening of his hearing. His deafness drove him to social isolation and his last public performance was at the age of forty-four³.

Antonio Valsalva in 1704 was the first person to describe the stapes ankylosis as a cause of hearing loss when describing bony obliteration of the oval window in a patient of acquired deafness³. In 1841, Toynebee described fixation of the stapes to the margins of the

oval window in 136 of 1000 temporal bone dissections. Politzer demonstrated the histology of otosclerosis and he identified the disease as an otic capsule disorder, characterized by abnormal new bone formation⁴.

In 1872, Anton von Trötsch coined the term otosclerosis, but did not make any difference between otosclerosis and tympanosclerosis⁵. Politzer, in 1894, recognized otosclerosis as the final stage of the disease with sclerotic plaques in the otic capsule⁶.

Development of Surgical Management of Otosclerosis Surgery for otosclerosis has developed through three distinct eras:

- 1) The mobilization era
- 2) The fenestration era
- 3) The stapedectomy era

The Mobilization Era

Prosper Meneire first reported mobilization of the stapes in 1842. In the late 1800s, Kessel attempted stapes mobilization without ossicular reconstruction in cases with ossicular fixation, thereby ushering in the mobilization era. Kessel described a case in the mid-1800's wherein hearing improvement was noted following a skull fracture in a young man with otosclerosis. Unfortunately the patient expired due to complications of head injury, and postmortem examination of his temporal bones revealed a fracture running through the horizontal semicircular canal. Based on this

observation, Kessel tried to create a similar fistula in patients with otosclerosis using a hammer and gauge and hence, he is considered to be the actual founder of stapes surgery⁷.

In 1891, Jackin Boston, USA, left the oval window open after removing the stapes. At the time, eminent contemporaries condemned the procedure due to its inherent risk off atal complications. The techniques introduced by Kessel and Jackdid not employ middle ear amplification structures, but allowed increased transmission of sound through the oval window. Also, any gains in hearing were frequently temporary because of refixation of any remaining stapes footplate. Fatal cases of meningitis were reported as occurring secondary to intraoperative exposure of perilymph to bacteria⁸.

Several French otolaryngologists performed mobilization of the stapes, including Boucheron and Miot. In 1890, Miot published a series of 200 cases of stapes mobilizations, applying the antiseptic principles of Lister with cleansing of the ear canal and sterilization of instruments. The results of Miot's stapediolysis were quite good, with hearing improvement in 59% of ears. Schwartze and Lucae from Germany⁹, Ferrari from Italy, Blakeio and Jack⁸ from USA and Sexton¹¹ and Alderton¹² from New York carried out stapes mobilization and removal of the stapes. Adam Politzer, Siebenmann and Moure, who were theleading otologists of their time, declared that stapes surgery was useless, dangerous and unethical at the 6th International Otology Congress in London⁹.

The Fenestration Era

Passov suggested the promontory fenestration in 1897, and in 1899 Floderus suggested an opening of the vestibular labyrinth. This was described by Jenkins in London in 1913 as a fenestration of the lateral semicircular canal. In the 1920s, Nylen in Sweden was the first to use a microscope for ear surgery¹³.

With the advent of the operating microscope, the fenestration era began in 1923, when Gunnar Holmgren (who is considered the Father of fenestration surgery) created a fistula in the lateral semicircular canal and sealed it immediately with periosteum¹⁴. This procedure worked by allowing preferential sound conduction via the fistula instead of the ossicular chain. Holmgren's procedure was popularized during the 1930's by Sourdille in France, who developed a three-stage technique which was widely published¹⁵. A one-stage technique for horizontal semicircular canal fenestration was developed by Julius Lempert in New York, which went on to achieve international recognition after it proved to improve hearing. However, the seresults were fleeting as the fenestra often resealed with bone and the technique was further refined by others¹⁶. These surgeries were aided by the development of improved anaesthesia, lighting, operating microscopes, antiseptic techniques and antibiotics. John House described the double blue line technique to avoid drilling directly over the fenestra.

The Stapedectomy Era

The stapedectomy era was ushered in prior to the end of the fenestration era. It was in 1952 when Samuel

Rosen from New York, tested the mobility of the stapes using a transcanal approach before a semicircular canal fenestration and hence, rediscovered that stapes mobilization causes hearing improvement¹⁷. John Shea Jr. later came to the conclusion that it must be possible to replace an otosclerotic stapes with a prosthesis. In collaboration with Treace, an engineer, he created a stapes prosthesis made of the biocompatible material Teflon. In a female patient with otosclerosis, after removal of the stapes and covering the oval window with a vein graft, he used this Teflon prosthesis for the first time on 1st May 1956 with complete success. After the Teflon stapes, Shea used a hollow polyethylene tube for a certain period of time, but this sometimes caused inner ear fistulae. Later, he used a piston made entirely of Teflon which is still used today by many surgeons¹⁸. In 1958, Shea advocated total stapes footplate removal when using the polyethylene strut prosthesis and a vein graft to seal the oval window. Shortly, thereafter Schuknecht advocated the use of wire and fat prosthesis¹⁹.

Portmann and Claverie in 1957 and Zangemeister in 1958 suggested that the suprastructure of the stapes should be left in situ and one of the stapes crura used as an interposition. The advantages of this is that no foreign material is implanted. In 1960, Zollner replaced the extracted stapes with an autologous cortical bone chip whereas Pflatz made a graft with a cartilage chip. In 1960, Schuknecht developed a steel wire-adipose tissue prosthesis made directly during the operation, which had the disadvantage of lateral displacements of the wire²⁰.

In the 1960s, Plester suggested the technique of partial stapedectomy in which only the posterior third of the foot plate was removed. In 1961, the piston concept was introduced in which a cup or piston prosthesis was used with connective tissue graft of vein to seal the oval window. The classic cup/piston prosthesis was fabricated of 316 L stainless steel, an alloy that is inert in living tissues and is non magnetic.²¹

In 1962, Shea et al and Marquet and Martin made a small opening in the middle of the footplate into which the prosthesis piston fitted exactly, in order to reduce the inner ear risk. This initiated the era of "stapedotomy" which has continued till the present time. Reverse Stapedotomy was popularized by Fisch and involved insertion of a prosthesis before removal of suprastructure of stapes. The advantage of this technique was that the ossicular chain remained very stable during insertion of prosthesis. Intact tendon stapedotomy preserves blood supply to lenticular and long process of incus, prevents slipping of the prosthesis and protects the inner ear against loud noise. House and Greenfield²² later advocated the use of wire with absorbable gelatin sponge (gelfoam) prosthesis. Use of wire/gelfoam or wire/fat prosthesis necessitated total removal of footplate for the wire prosthesis to function satisfactorily. Moon²³ advocated cup/piston prosthesis with areolar tissue. Glasscock²⁴ and Sheehy²⁵ and their associates Lippy and Schuring^{26,27} advocated leaving the oval window neomembrane intact and undisturbed, if possible, in revision cases, to reduce the risk of severe SNHL. Feldman and Schuknecht²⁸, Pearman and Dawes²⁹, and Derlacki³⁰ reported opening the

neomembrane to identify the vestibule and ensure correct prosthesis placement.

Newer Advances And The Future Of Stapes Surgery

Perkins in 1980 performed laser surgery in otosclerosis for the first time. Laser Stapedotomy provides pinpoint accuracy and instantaneous cauterization minimising tissue trauma. The Argon laser has been more commonly used in stapedotomy because it has a visible wavelength and better accuracy which helps to prevent inner ear damage and also causes photo coagulation. Poe in 2000 advocated for laser-assisted endoscopic stapedectomy, which provides a better view of the surgery³¹.

Laser stapedectomy minus prosthesis (LASER STAMP) is a physiological stapedectomy done in early cases of Otosclerosis involving the fissula ante fenestram. In this procedure, the anterior crus is vapourised using Argon laser and the anterior third of the otosclerotic footplate is separated from the remaining stapes. Here, the posterior part of the footplate with the posterior crus and the incudostapedial joint, acts like a piston³². Most recently, Rothbaum et al from John Hopkins, Baltimore, have published a report on robot-assisted stapedotomy, specifically concerning the use of robot ics for micropick fenestration of the stapes surgery³³.

Conclusion

Stapes surgery has come a long way since its inception in the 1800s thanks to the efforts of the pioneers mentioned above, and many of the techniques developed by them persist today, albeit with many variations. However, the fundamental principle of re-establishing a mobile link between the tympanic membrane and vestibule has remained the objective of surgery for patients with otosclerosis. Although certain challenges linger, such as patients of otosclerosis with sensorineural hearing loss and vertigo, for now, the future looks bright for further development of surgery of the stapes.

The authors declare no conflict of interest.

References

- 1) Velegrakis GA. Otosclerosis: state of the art. *Otorhinolaryngologia Head Neck Surg* 2011; 43:6-16.
- 2) Khan H, Shah JI, Saeedullah, Khan M, Asad K. Small fenestra stapedotomy. *JPML*;18(2):176-181.
- 3) Valsalva AM: The Human Ear (Latin) Bononiae: C.Pisari, 1704
- 4) Toynbee J. Pathological and surgical observations of the diseases of the ear. *Trans Med Chir Soc Lond*. 1841;24:190-196.
- 5) Welling DB, Glasscock ME, Gantz BJ. Avulsion of the anomalous facial nerve at stapedectomy. *Laryngoscope*. 1992; 102: 729-733.
- 6) Roland PS, Meyerhoff WL. Otosclerosis. In: Bailey BJ, Calhoun KH (Hrsg) *Head and neck Surgery – Otolaryngology*. Lippincott Williams and Wilkins, New York, 2001; S1831-1841.
- 7) Arnold W, Hausler R (eds): *Otosclerosis and Stapes Surgery*. *AdvOtorhinolaryngol*, Basel, Karger, 2007;65:1-5.
- 8) Jack FL. Remarkable improvement of hearing by removal of the stapes. *Trans Am Otol Soc*. 1893;284:474-89.
- 9) Arnold W, Hausler R (eds): *Otosclerosis and Stapes Surgery*. *AdvOtorhinolaryngol*, Basel, Karger, 2007;65:1-5.
- 10) Blake CJ: Operation for removal of the stapes. *Box M & S J* 1892;127:469.
- 11) Sexton S: Operation for deafness and tinnitus, due to immobilization of the ossicles, and for otorrhea. *Br Med J* 1890;1:69-70.
- 12) Alderton HA: Trephining of the stapedial footplate for otitis media sclerososa. *Trans Am Otol Soc* 1898;3:60-63.
- 13) Jenkins GJ: Otosclerosis: Certain clinical features and experimental operative procedures. *Trans XVIIth Inter Med London* 1913;16:609-618.
- 14) Holmgren J. The surgery of otosclerosis. *Ann OtolRhinolLaryngol*. 1937;46:3-12.
- 15) Sourdille M. New technique in the surgical treatment of severe and progressive deafness from otosclerosis. *Bull NY Acad Med*. 1937;13:673.
- 16) Lempert J. Improvement in hearing in cases of otosclerosis: A new, one-stage surgical technic. *Arch Otolaryngol*. 1938;28:42-97.
- 17) Rosen S. Restoration of hearing in otosclerosis by mobilization of the fixed stapedial footplate; an analysis of results. *Laryngoscope*. Apr 1955;65(4):224-69.
- 18) Shea JJ Jr. A personal history of stapedectomy. *Am J Otol*. Sep 1998;19(5 Suppl):S2-12.
- 19) Shea JJ Jr : Fenestration of the oval window. *Ann otolrhinolaryngol* 1958;57:932.
- 20) Schuknecht H :stapedectomy and graft prosthesis operation. *Actaotolaryngol(stockh)* 1960;51: 241-243
- 21) Robinson M: The stainless-steel stapedial prosthesis: one year's experience. *Laryngoscope* 1962;73:514
- 22) House HP, Greenfield EC : Five-year study of wire loop absorbable gelatin sponge technique. *Arch otolaryngol Head Neck surg* 1969;89:420-421.
- 23) Moon C: Stapedectomy connective tissue graft and the stainless-steel prosthesis. *Laryngoscope* 1968;78:798-807.

- 24) Glasscock ME III, Mckennan KX, Levine SC: Revision stapedectomy surgery. *Otolaryngol Head neck surg* 1987;96:141-148.
- 25) Sheely JL, Nelson RA, House HP: Revision stapedectomy: A review of 258 cases. *Laryngoscope* 1981;91:43-51
- 26) Lippy WH, Schuring AG: Stapedectomy revision of the wire gelfoam prosthesis. *Otolaryngol Head neck surg* 1983;91:9-13
- 27) Lippy WH, Schuring AG: Stapedectomy revision following sensorineural hearing loss. *Otolaryngol Head neck surg* 1984;92:580-582
- 28) Feldman BA, Schuknecht HF: Experiences with revision stapedectomy procedures. *Laryngoscope* 1970;80:1281-1291
- 29) Pearman K, Dawes JDK: Post stapedectomy conductive deafness and results of revision surgery. *J Laryngol Otol* 1982;96:405-410
- 30) Derlacki EL: Revision stapes surgery: problems with same solutions. *Laryngoscope* 1985;95:1047-1053
- 31) Poe DS. Laser-assisted endoscopic stapedectomy: a prospective study. *Laryngoscope*. May 2000;110(5 Pt 2 Suppl 95):1-37.
- 32) Silverstein H. Laser stapedotomy minus prosthesis (laser STAMP): a minimally invasive procedure. *Am J Otol*. May 1998;19(3):277-82.
- 33) Rothbaum DL et al. Robot-assisted stapedotomy: micropick fenestration of the stapes footplate. *Otolaryngol Head Neck Surg* 2002;127(5):417-26.

Fasting Unnecessary for Lipid Profile

Excluding Denmark, all other countries of the world require the patients to fast for at least 8 hours before undergoing tests for cholesterol and other components of lipid profile. In Denmark, non-fasting samples have been in use for lipid profile since 2009. The problem with fasting samples is one of compliance, particularly from children, senior citizens, diabetic patients and patients on certain drug therapies. Besides, Insistence on fasting makes it difficult to do population screening. Now an expert group representing the European Atherosclerosis society (EAS) and the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM) have recommended that non-fasting samples are adequate for majority of the patients for the estimation of cholesterol and other lipid components. This recommendation comes after careful evaluation of research done in USA, Canada and Denmark covering well over 300,000 study subjects. However, fasting sample is required when triglyceride levels are more than 440/dL (5 mmol/L) (*European Heart Journal*, doi: 10. 1093/eurheartj/ehw152, published online 26 April 2016).

- Dr. K. Ramesh Rao